

-1-

HUMAN NEURONAL NICOTINIC ACETYLCHOLINE RECEPTOR
COMPOSITIONS AND METHODS EMPLOYING SAME

Field of the Invention

5

This invention relates to neuronal nicotinic acetylcholine receptor genes and proteins. In a particular aspect, the present invention relates to human neuronal nicotinic acetylcholine receptor genes and proteins. In a further aspect, the present invention relates to methods for determining the presence of neuronal nicotinic acetylcholine receptor activity in cells thought to have genes encoding such proteins. In yet another aspect, the present invention relates to methods for determining the agonist or antagonist activity of compounds which might interact with neuronal nicotinic acetylcholine receptors.

10

15

Background of the Invention

20

Most theories on how the nervous system functions depend heavily on the existence and properties of cell to cell contacts known as synapses. For this reason, the study of synapses has been a focal point for neuroscience research for many decades.

25

Because of its accessibility to biochemical and electrophysiological techniques, and because of its elegant, well defined structure, the neuromuscular synapse (also known as the neuromuscular junction), which occurs at the point of nerve to muscle contact, is one of the most studied and best understood synapses. At the neuromuscular junction, the nerve cell releases a chemical neurotransmitter, acetylcholine, which binds to nicotinic acetylcholine receptor proteins located on post-synaptic muscle.

30
35

-2-

cells. The binding of acetylcholine results in a conformational change in the nicotinic acetylcholine receptor protein. This change is manifested by the opening of a transmembrane channel in the receptor
5 which is permeable to cations. The resulting influx of cations depolarizes the muscle and ultimately leads to muscle contraction.

Biological and structural studies have shown that the nicotinic acetylcholine receptor in muscle is
10 a glycoprotein composed of five subunits with the stoichiometry $\alpha\alpha\beta\gamma\delta$ (alpha-alpha-beta-gamma-delta). From these same studies, it is known that each of the subunits has a mass of about 50-60 kilodaltons and is encoded by a separate gene. In vitro reconstitution
15 experiments have shown that this $\alpha\alpha\beta\gamma\delta$ complex is a functional receptor containing both ligand binding sites and a ligand-gated transmembrane channel.

It is now known that a variety of neurotransmitters and neurotransmitter receptors exist
20 in the central and peripheral nervous systems. Despite this knowledge, there is still little understanding of the diversity of receptors for a particular neurotransmitter, or of how this diversity might generate different responses to a given
25 neurotransmitter, or to other modulating ligands, in different regions of the brain. On a larger scale, there is little appreciation of how the use of a particular synapse makes it more or less efficient, or
hnges in neuronal circuits might be accomplished by the
30 modification of synapses.

An understanding of the molecular mechanisms involved in neurotransmission in the central nervous system is limited by the complexity of the system. The cells are small, have extensive processes, and often
35 have thousands of synapses deriving from inputs from

3

-3-

many different parts of the brain. In addition, the actual number of neurotransmitter receptors is low, making their purification difficult, even under the best of circumstances. Consequently, neither cellular
5 nor biochemical approaches to studying neurotransmission in the central nervous system has been particularly fruitful. This is unfortunate because it is quite probable that the treatment of dementia, Alzheimer's disease and other forms of mental
10 illness will involve modification of synaptic transmission with specific drugs.

Nicotinic acetylcholine receptors found at the vertebrate neuromuscular junction, in vertebrate sympathetic ganglia and in the vertebrate central
15 nervous system can be distinguished pharmacologically on the basis of ligands that open or block the ion channel. For example, the elapid α -neurotoxins that block activation of nicotinic acetylcholine receptors at the neuromuscular junction do not block activation
20 of neuronal nicotinic acetylcholine receptors found on several different cell lines.

To gain access to the neuronal acetylcholine receptors, traditional biochemical and neurophysiological methods have been abandoned in favor
25 of the newer methods of molecular biology. More specifically, using molecular cloning techniques, complementary DNA clones were isolated which encode the acetylcholine receptor expressed in the Torpedo fish electric organ, a highly enriched source of receptor.
30 The cDNA clones isolated from the fish electric organ were then used in nucleic acid hybridization experiments to obtain cDNA and genomic clones for the subunits of the acetylcholine receptor expressed in mouse skeletal muscle.

4

-4-

The availability of cDNA clones encoding muscle nicotinic receptors made it possible to extend these studies in the important direction of neuronal receptors. More specifically, based on the assumption
5 that neuronal nicotinic receptors are evolutionarily related to muscle receptors, and that this relationship will be reflected at the genetic level by nucleotide sequence homology, the cDNA clones encoding the muscle nicotinic receptor were used to screen rat cDNA and
10 genomic libraries for related neuronal mRNAs or genes. This method has resulted in the isolation of several neuronal cDNA clones that have significant sequence homology with the muscle acetylcholine clones.

That the neuronal nicotinic acetylcholine
15 receptors differ from muscle nicotinic acetylcholine receptors is evidenced by the fact that neuronal receptors can be constituted from only two different gene products (i.e., one alpha subunit and one beta subunit). This is significant since, in all
20 experiments reported to date, muscle nicotinic acetylcholine receptors have been formed with $\alpha\beta\delta\Delta$ subunits, $\alpha\beta\Delta$ subunits, $\alpha\beta\delta$ subunits or $\alpha\delta\Delta$ subunits, but not with any pairwise combinations. See Kurosaki et al., FEBS Letters 214, 253-258 (1987).

25 In order to further extend such studies, to provide proteins useful for assaying compounds as potential agonists or antagonists for human neuronal nicotinic acetylcholine receptors, as well as cell lines capable of expressing such proteins, we undertook
30 to isolate and characterize clones which encode various subunits of the human neuronal nicotinic acetylcholine receptor; we further undertook to develop methods for expressing cloned human neuronal nicotinic acetylcholine receptor sequences in recombinant cell
35 lines; and we further undertook to develop assays for

5

-5-

identifying which of the resultant recombinant cell lines express functional neuronal nicotinic receptors.

Summary of the Invention

5

In accordance with the present invention, we have isolated and characterized clones which encode the alpha2, alpha3 and beta2 subunits of the human neuronal nicotinic acetylcholine receptor.

10

The neuronal clones of the present invention encode a family of acetylcholine receptors having unique pharmacological properties. The demonstration that the nicotinic acetylcholine receptors are much more diverse than previously expected offers an opportunity for a high level of pharmaceutical intervention and a chance to design new drugs that affect specific receptor subunits. Such subtypes make it possible to observe the effect of a drug substance on a particular receptor subtype, which can be expressed in a recombinant cell in the absence of the other receptor subtypes. Information derived from these observations will allow the development of new drugs that are more specific, and therefore have fewer unwanted side effects.

25

In addition, the availability of human neuronal receptors makes it possible to perform initial in vitro screening of the drug substance in a test system which is specific for humans. While it is true that the drug eventually has to be administered directly to the human patient, it is probable that useful drugs are being missed because conventional drug screening is limited to assays employing non-human receptors, human tissue preparations (which are likely to be contaminated with other receptors, both nicotinic and non-nicotinic in origin), and other suboptimal

35



-6-

assay systems. Consequently, the ability to screen drug substances in vitro on specific receptor subtype(s) is likely to be more informative than merely screening the drug substance employing presently available suboptimal assay systems.

Both the receptor subunit genes and proteins of the present invention can be used for drug design and screening. For example, the cDNA clones encoding the human alpha2, alpha3 and beta2 receptor subunits can be transcribed in vitro to produce mRNA. This mRNA, either from a single subunit clone or from a combination of clones, can then be injected into oocytes where the mRNA directs the synthesis of the human receptor molecule(s). The resulting receptor-expressing oocytes can then be contacted with a test compound, and the agonist or antagonist effect thereof can then be evaluated by comparing oocyte response relative to positive and negative control compounds and positive and negative control oocytes. Alternatively, the clones may be placed downstream from appropriate gene regulatory elements and inserted into the genome of eukaryotic cells. This will result in transformed cell lines expressing a specific human receptor subtype, or specific combinations of subtypes. The derived cell lines can then be produced in quantity for similar reproducible quantitative analysis of the effects of drugs on receptor function.

Brief Description of the Figures

Figure 1 is a restriction map of the alpha2 subunit gene of the human neuronal nicotinic acetylcholine receptor, compared to the corresponding rat gene.

7

-7-

Figure 2 is a restriction map of the alpha3 subunit gene of the human neuronal nicotinic acetylcholine receptor, compared to the corresponding rat gene.

5 Figure 3 is a restriction map of the beta2 subunit gene of the human neuronal nicotinic acetylcholine receptor, compared to the rat gene.

I
10 Figure 4^{A-C} is a comparison of the alpha2 subunit gene of the rat neuronal nicotinic acetylcholine receptor with several cDNA fragments obtained from the human alpha2 subunit gene. The arrows beneath the various human cDNA fragments indicate the direction and extent of DNA sequencing carried out for the respective fragments.

I
15 Figure 5^{A-B} is a comparison of the alpha3 subunit gene of the rat neuronal nicotinic acetylcholine receptor with several cDNA fragments obtained from the human alpha3 subunit gene. The arrows beneath the various human cDNA fragments
20 indicate the direction and extent of DNA sequencing carried out for the respective fragments.

I
25 Figure 6^{A-E} is a comparison of the beta2 subunit gene of the rat neuronal nicotinic acetylcholine receptor with several cDNA fragments obtained from the human beta2 subunit gene. The arrows beneath the various human cDNA fragments indicate the direction and extent of DNA sequencing carried out for the respective fragments.

I
30 Figure 7^{A-B} is a comparison of about 500 base pairs of human alpha2 sequence with the corresponding rat sequence.

I
 Figure 8^{A-B} is a comparison of about 650 base pairs of human alpha3 sequence with the corresponding rat sequence.

8

-8-

T
A-E
Figure 9 is a comparison of the nucleotide sequence for the human and rat beta2 subunits.

Detailed Description of the Invention

5

In accordance with the present invention, there are provided substantially pure DNA sequence(s) encoding alpha subunit(s) of the human neuronal nicotinic acetylcholine receptor and/or substantially
10 pure DNA sequence(s) encoding beta subunit(s) of the human neuronal nicotinic acetylcholine receptor.

In accordance with a particular embodiment of the present invention, there are provided mRNA sequences and polypeptides encoded by the above-
15 described DNA sequences.

In accordance with yet another embodiment of the present invention, there are provided cells transformed with one or more of the above-described DNA sequences.

20 In accordance with still another embodiment of the present invention, there are provided substantially pure human neuronal acetylcholine receptors comprising at least one human alpha receptor subunit and at least one human beta subunit.

25 In accordance with a further embodiment of the present invention, there are provided methods for measuring the agonist or antagonist activity of test compounds (with respect to human neuronal acetylcholine receptors or subunits thereof), by measuring the
30 response of the above-described cells and/or receptors, relative to the response of a control, when contacted with said compound.

In accordance with the latter embodiment of the present invention, the response of the above-

9

-9-

described cells and/or receptors is determined by such assays as:

5 nicotine binding,
 ^{86}Rb ion-flux,
 the electrophysiological response of
said cells, or

 the electrophysiological response of
oocytes transfected with RNA from said cells.

10 In accordance with yet another embodiment of
the present invention, there is provided a method for
assaying cells for the presence of neuronal nicotinic
acetylcholine receptor activity. This is accomplished
by determining the effect of known neuronal nicotinic
acetylcholine agonists and/or antagonists on the influx
15 of ^{86}Rb ions into cells, relative to the rate of influx
of ^{86}Rb ions into control cells.

 In accordance with a further embodiment of
the present invention, there is provided an alternative
method for assaying cells for the presence of neuronal
20 nicotinic acetylcholine receptor activity, employing a
multi-step screening protocol comprising the steps:

 (a) analyzing said cells for the
presence of alpha and beta subunit RNAs,

25 (b) analyzing those cells which are
positive for the presence of alpha and beta subunit
RNAs for their ability to bind nicotine or a nicotine
agonists, relative to the nicotine binding ability of
control cells known to express neuronal nicotinic
acetylcholine receptors, and

30 (b) determining the effect of known
neuronal nicotinic acetylcholine agonists and/or
antagonists on cells having the ability to bind
nicotine or nicotine agonist on the influx of ^{86}Rb ions
into said cells, relative to the rate of influx of ^{86}Rb
35 ions into control cells.

10

-10-

In accordance with a still further embodiment of the present invention, there is provided a method for making cells having neuronal nicotinic acetylcholine receptor activity, employing a multi-step protocol comprising the steps:

(a) transfecting host cells with DNA encoding at least one alpha subunit of the neuronal nicotinic acetylcholine receptor and at least one beta subunit of the neuronal nicotinic acetylcholine receptor, then

(b) analyzing said transfected cells for the presence of alpha and beta subunit RNAs, employing methods such as Northern blot or slot blot analysis; then

(c) analyzing those cells which are positive for the presence of alpha and beta subunit RNAs for their ability to bind nicotine or a nicotine agonist, relative to the nicotine binding ability of control cells known to express neuronal nicotinic acetylcholine receptors, and

(d) determining the effect of known neuronal nicotinic acetylcholine agonists and/or antagonists on cells having the ability to bind nicotine or a nicotine agonist on the influx of ⁸⁶Rb ions into control cells.

In accordance with the preceding two embodiments of the present invention, mRNA from cells which are positive for alpha and beta neuronal nicotinic acetylcholine subunits is injected into oocytes, which are then assayed for the presence of functional neuronal nicotinic acetylcholine receptors.

As used herein, the term agonist refers to a substance that activates receptor function; and the term antagonist refers to a substance that interferes with receptor function. Antagonists are of two types:

-11-

competitive and non-competitive. A competitive antagonist (or competitive blocker) competes with the neurotransmitter for the same binding site. A non-competitive antagonist or blocker inactivates the functioning of the receptor by binding to a site other than the acetylcholine binding site.

As used herein, alpha2 refers to a gene, which has been identified in chick and rat, that encodes a neuronal subunit of the same name. DNA coding for the human neuronal alpha2 subunit has been deposited with the ATCC; the DNA (designated as HnAChR α 2; a restriction map of which is shown in Figure 1; and a partial nucleotide sequence of which is shown in Figure 7) has been accorded ATCC Accession No. 68277.

As used herein, alpha3 refers to a gene that encodes a neuronal subunit of the same name. DNA coding for the human neuronal alpha3 subunit has been deposited with the ATCC; the DNA (designated as HnAChR α 3; a restriction map of which is shown in Figure 2; and a partial nucleotide sequence of which is shown in Figure 8) has been accorded ATCC Accession No. 68278.

As used herein, beta2 refers to a gene encoding a neuronal nicotinic acetylcholine subunit of the same name. DNA coding for the human neuronal beta2 subunit has been deposited with the ATCC; the DNA (designated as HnAChR β 2; a restriction map of which is shown in Figure 3; and the nucleotide sequence of which is shown in Figure 9) has been accorded ATCC Accession No. 68279.

cDNA clones comprising human neuronal nicotinic acetylcholine receptor genes alpha2 (clone HnAChR α 2), alpha3 (clone HnAChR α 3), and beta2 (clone HnAChR β 2), all of which are in E. coli HB101, have been

12

-12-

deposited at the American Type Culture Collection,
Rockville, Maryland, U.S.A. (ATCC) under the terms of
the Budapest Treaty on the International Recognition of
Deposits of Microorganisms for Purposes of Patent
5 Procedure and the Regulations promulgated under this
Treaty. Samples of the cloned genes are and will be
available to industrial property offices and other
persons legally entitled to receive them under the
terms of said Treaty and Regulations and otherwise in
10 compliance with the patent laws and regulations of the
United States of America and all other nations or
international organizations in which this application,
or an application claiming priority of this
application, is filed or in which any patent granted on
15 any such application is granted.

Use of the phrase "substantial sequence
homology" in the present specification and claims means
that DNA, RNA or amino acid sequences which have slight
and non-consequential sequence variations form the
20 actual sequences disclosed and claimed herein are
considered to be equivalent to the sequences of the
present invention, and as such are within the scope of
the appended claims. In this regard, "slight and non-
consequential sequence variations" mean that
25 "homologous" sequences (i.e., the sequences that have
substantial sequence homology with the DNA, RNA, or
proteins disclosed and claimed herein) will be
functionally equivalent to the sequences disclosed and
claimed in the present invention. Functionally
30 equivalent sequences will function in substantially the
same manner to produce substantially the same
compositions as the nucleic acid and amino acid
compositions disclosed and claimed herein.

Use of the phrase "substantially pure" in the
35 present specification and claims as a modifier of DNA,

B

-13-

RNA, polypeptides or proteins means that the DNA, RNA, polypeptides or proteins so designated have been separated from their in vivo cellular environments through the efforts of human beings; as a result of this separation, the substantially pure DNAs, RNAs, polypeptides and proteins are useful in ways that the non-separated, impure DNAs, RNAs, polypeptides or proteins are not.

The invention DNA sequences were isolated employing analogous rat neuronal acetylcholine receptor subunit DNA fragments as probes in various human cDNA libraries. Due to the very low concentration of various human neuronal subunits in their native state, the frequently very localized presence of some of the human neuronal subunits in various sources of tissue, the difficulty in obtaining human neuronal (brain) tissue with which to work, as well as the high level of care necessary to ensure the presence of intact mRNA in the source human neuronal tissue, a significant problem to be solved in order to achieve the objects of the present invention was identifying and obtaining suitable source(s) of DNA to probe for the desired sequences. By probing numerous human cDNA libraries, e.g., pre-frontal cortex cDNA, parietal cDNA, temporal cortex cDNA, brain stem cDNA, basal ganglia cDNA, and spinal cord cDNA, various fragments of the human neuronal subunits were identified (see, for example, Figures 4^{A-C}, 5^{A-B} and 6^{A-E}). After partial sequencing and restriction mapping of several such fragments, and comparison of such fragments to the analogous rat sequences, it was possible to identify composite DNA sequences for the human neuronal alpha2, alpha3 and beta2 subunits, as disclosed and claimed herein.

In addition to their use as coding sequences for the production of human neuronal subunits and

14

-14-

synthetic human neuronal receptors, the invention sequences can also be used as probes for the identification of additional human neuronal sequences. This is done by probing various sources of human neuronal DNA with invention sequences, then selecting those sequences having a significant level of sequence homology with the probe employed.

Invention DNA sequences can be transformed into a variety of host cells. Eukaryotic cells such as yeast or mammalian cells are presently preferred. A variety of suitable host mammalian cells, having desirable growth and handling properties, are readily available to those of skill in the art. Especially preferred for such purpose are human, rat or mouse cells.

Similarly, a variety of suitable yeast cells are readily available to host cells for the invention sequences. Especially preferred are yeast selected from Pichia pastoris, Saccharomyces cerevisiae, Candida tropicalis, Hansenula polymorpha, and the like.

Alternatively, the invention DNA sequences can be ^{transcribed} translated into RNA, which can then be transfected into amphibian cells for ^{translation} transcription into protein. Suitable amphibian cells include Xenopus oocytes.

Cells transformed with invention DNA (or RNA) can optionally be further transformed with a reporter gene expression construct, so as to provide a ready, indirect measure of the presence of functional human neuronal receptor in the transformed cell. Such a reporter gene expression construct comprises:

a transcriptional control element; wherein said transcription control element, in said cell, is responsive to an intracellular condition that occurs when the human neuronal nicotinic acetylcholine

15

-15-

receptor interacts with a compound having agonist or antagonist activity with respect to said receptor, and
a reporter gene encoding a transcription and/or translational product; wherein said product can
5 be, directly or indirectly, readily measured; and
wherein said gene is in operative association with said transcriptional control element.

Transcriptional control elements contemplated for use in this embodiment of the present invention
10 include the c-fos promoter, the vasoactive intestinal peptide gene promoter, the somatostatin gene promoter, the proenkephalin gene promoter, the phosphoenolpyruvate carboxykinase gene promoter, the NGFI-A gene promoter, and the like.

15 Reporter genes contemplated for use in this embodiment of the present invention include the chloramphenicol transferase (CAT) gene, the gene product of which can be readily analyzed by a variety of methods known in the art. See, for example,
20 Nielsen, et al., Anal. Biochem. 179, 19-23 (1989), luciferase and other enzyme detection systems such as alkaline phosphatase, β -galactosidase, and the like.

A particularly useful application of the
25 invention sequences is the ability to prepare synthetic receptors and synthetic receptor subunits which are substantially free of contamination from other, potentially competing proteins. Thus, a cell transformed with the invention alpha2 and beta2
30 sequences could express a synthetic receptor consisting essentially of only the alpha2 and beta2 subunits. Such a synthetic receptor would be useful for a variety of applications, e.g., as part of an assay system free of the interferences frequently present in prior art

16

-16-

assay systems employing non-human receptors or human tissue preparations.

Similarly, a synthetic receptor could be prepared by causing cells transformed with the invention alpha3 and beta2 sequences to express the corresponding proteins. The resulting synthetic receptor would consist essentially of only the alpha3 and beta2 subunits. Such a synthetic receptor would be useful for a variety of applications, e.g., as part of an assay system free of the interferences frequently present in prior art assay systems employing non-human receptors or human tissue preparations.

Furthermore, testing of single receptor subunits with a variety of potential agonists or antagonists would provide additional information with respect to the function and activity of the individual subunits. Such information may lead to the identification of compounds which are capable of very specific interaction with one or more of the receptor subunits. Such specificity may prove of great value in medical application.

In accordance with one aspect of the present invention, assay methods have been developed for the ready determination of the presence of functional neuronal nicotinic acetylcholine receptors. Thus, cells transformed with invention DNA or RNA sequences, or cell-lines derived from a variety of other sources can be readily screened to determine if functional receptors are produced thereby. One useful assay method is the "⁸⁶Rb ion-flux" assay, wherein the influx of ⁸⁶Rb ions into test cells is measured as a function of the presence or absence of known neuronal nicotinic acetylcholine agonists or antagonists. Thus, a cell which shows no difference in the ⁸⁶Rb ion flux, whether in the presence or absence of agonist or antagonist is

17

-17-

not expressing functional neuronal receptor. This assay provides more information than is provided by a simple binding assay because it also indicates whether or not functional receptor is present.

5 Another useful assay method of the invention involves subjecting test cells to the following steps:

(a) analyzing said cells for the presence of alpha and beta subunits RNAs,

10 (b) analyzing those cells which are positive for the presence of alpha and beta subunit RNAs for their ability to bind nicotine or a nicotine agonist, relative to the nicotine binding ability of control cells known to produce neuronal nicotinic acetylcholine receptors, and

15 (c) determining the effect of known neuronal nicotinic acetylcholine agonists and/or antagonists on cells having the ability to bind nicotine or nicotine agonist on the influx of ^{86}Rb ions into said cells, relative to the rate of influx of ^{86}Rb ions into
20 positive and/or negative control cells.

Cells can be analyzed for the presence of alpha and beta subunit RNA in a variety of ways, such as for example, by Northern hybridization, slot blot analysis, and the like.

25 The determination of the nicotine-binding ability of test cells can readily be determined by techniques known by those of skill in the art. For additional detail, see Example 3B below.

The ^{86}Rb ion-flux assay is then carried out as
30 described hereinabove.

The above-described sequence of analytical steps provides an effective way to screen large numbers of transformed cells for the expression of neuronal receptor subunit(s), the ability of such subunit(s) to
35 bind to nicotine, nicotine agonists or nicotine

18

-18-

antagonists, and the ability of such subunit(s) to assemble into functional receptors.

As a further step to verify the ability of test cells to produce functional receptor, mRNA from
5 cells which are positive for the presence of alpha and beta neuronal nicotinic acetylcholine receptor subunits by the above-described assays can be injected into oocytes, which can then be assayed for the presence of functional neuronal nicotinic acetylcholine receptors.
10 As another alternative, one can measure the electrophysiology of the positive cells (either directly or upon expression of RNA by oocytes). Positive results in each of these assays provides one with a high level of confidence that the test cells
15 contain the coding sequences for the production of receptor, and that such receptor is indeed expressed.

In accordance with another aspect of the present invention, a method for making eukaryotic cells having neuronal nicotinic acetylcholine receptor
20 activity is provided. Eukaryotic cells (e.g., mammalian or yeast cells) are transfected with DNA encoding at least one alpha subunit and at least one beta subunit of the neuronal nicotinic acetylcholine receptor. The resulting cells are then screened by one
25 or more of the above-described assay methods to identify those cells which have successfully incorporated the desired DNA sequences.

The invention will now be described in greater detail by reference to the following non-
30 limiting examples.

19

-19-

EXAMPLES

Plasmids comprising insert DNA encoding human α and β subunit proteins of the neuronal nicotinic acetylcholine receptor (nNACHR), isolated from various sources of human neuronal tissue, have been deposited in the ATCC. The clone names and deposit numbers are:

<u>Subunit</u>	<u>Clone Name</u>	<u>ATCC Accession #</u>
$\alpha 2$	HnACHR $\alpha 2$	68277
$\alpha 3$	HnACHR $\alpha 3$	68278
$\beta 2$	HnACHR $\beta 2$	68279

Restriction maps of the nNACHR-encoding inserts in these clones, as compared to the corresponding rat cDNA maps, are provided in Figures 1, 2, and 3, respectively. The rat cDNA inserts are described in Wada *et al.* (1988), *Science* 240:330-334 ($\alpha 2$); Boulter *et al.* (1986), *Nature* 319:368-374 ($\alpha 3$); Boulter *et al.* (1987), *Proc. Natl. Acad. Sci.* 84:7763-7767 ($\beta 2$). EcoRI adapters (from cloning vector) are present on the ends of each insert.

Portions of the $\alpha 2$ and $\alpha 3$, and all of the $\beta 2$, human neuronal NACHR subunit-encoding sequence were sequenced. The sequences of the human cDNAs were compared to the corresponding regions of the rat cDNAs, and the percent homology between the human and rat sequences are provided in Figures 4, 5, and 6, respectively. The nucleotide sequence homology is presented outside the parentheses, the translated amino acid sequence homology is presented in parentheses.

Additionally, actual nucleotide sequence comparisons are presented in Figures 7, 8, and 9. In all figures the human sequence is on top and the rat sequence is on the bottom. The nucleotide numbers for

20

-20-

the rat sequences correspond to the actual nucleotide positions in the coding sequence. In contrast, the nucleotide numbers for the human sequences do not correspond to the coding sequences; instead, these
5 numbers are related to the individual sequenced fragments. Figure 7, ^{A-B} presents the nucleotide sequence of the human $\alpha 2$ sequence as compared to the rat $\alpha 2$ sequence starting around the common BamHI site (see Figure 4) and continuing in the 3' direction
10 approximately 500 nucleotides. The degree of nucleotide homology in section A of Figure 7 is 87% and in section B is 93%.

^{A-B} Figure 8 presents the nucleotide sequence of the human $\alpha 3$ sequence starting about 50 nucleotide 3' from the 5' end of the coding sequence, and continuing
15 in the 3' direction for about 650 nucleotides. The degree of nucleotide homology between the human and rat sequence in Section A is 86% and in Section B is 90%.

^{A-E} Figure 9 presents the entire coding sequence
20 of the cDNA encoding the human $\beta 2$ subunit. It has 87% homology to the rat sequence at the nucleotide level.

Example 1

25 CONSTRUCTION OF EUKARYOTIC EXPRESSION VECTORS COMPRISED OF THE HUMAN NEURONAL NACHR SUBUNIT SEQUENCES

The cDNAs encoding the human neuronal NACHR subunits were inserted into the eukaryotic expression vector pSV2+Ldhfr, the construction of which is
30 described in Example 2. Each insert was excised from its plasmid (HnAchR $\alpha 2$, HnAchR $\alpha 3$, or HnAch $\beta 2$) by digestion with EcoRI. The resultant fragments were gel purified and the ~2600 bp ($\alpha 2$), ~3200 bp ($\alpha 3$), and ~2450 bp ($\beta 2$) fragments were isolated. Each insert
35 fragment was ligated to EcoRI-digested and

21

-21-

dephosphorylated pSV2+Ldhfr; 0.1 μ g of each DNA was used. The ligation reaction was transformed into MC1061 cells and amp^R colonies were selected. The desired plasmid(s) having insert in the correct orientation was (were) identified by the diagnostic fragments provided below, and named as follows:

T220X

<u>Subunit</u>	<u>Plasmid name</u>	<u>Diagnostic fragment</u>
α 2	h α 2/pSV2	PvuII: 550, 100, 7000 bp
α 3	h α 3/pSV2	PvuII: 850, 7350 bp
β 2	h β 2/pSV2	HindIII: 450, 7000 bp

These plasmids have the subunit-encoding insert placed in functional association downstream of the SV40 early promoter.

Example 2

DEVELOPMENT OF MAMMALIAN CELL LINES EXPRESSING α AND β SUBTYPES OF THE RAT NEURONAL NICOTINIC ACETYLCHOLINE RECEPTOR (rNACHR)

Cell lines were developed in Ltk⁻ (mouse fibroblast thymidine kinase deficient) cells by cotransfecting a plasmid comprised of an α -subunit-encoding sequence, a plasmid comprised of a β -subunit-encoding sequence, and a plasmid comprised of either the wild-type or crippled TK gene. A reporter gene expression construct can also be cotransfected into the cells to provide a transcription-based assay system. While the following examples employ eukaryotic expression vectors comprised of the rat NACHR subunit cDNA sequences, the eukaryotic expression vectors comprised of the human nNACHR cDNA sequences (Example 1) also can be used.

35

22

-22-

A. Host Cells

Ltk⁺ cells are available from ATCC (accession #CCL1.3).

5

B. rNACHR α - and β -Expression Plasmids

The α - and β -encoding eukaryotic expression plasmids were constructed using a slightly modified pSV2dhfr parent plasmid [Subramani, et al. (1981). Mol. Cell. Biol. 1:854-864] and α - and β -encoding inserts from the rat nNACHR subunit clones. The clone sources for the subunit sequences were:

Subunit	Parent plasmid	Insert fragment
α 2	HYP16(9) ^a	~2 Kb <u>EcoRI</u>
α 3	PCA48E(4) ^b	~2 Kb <u>HindIII-EcoRI</u>
* α 4.1	HYA23-1E(1) ^c	~2 Kb <u>HindIII</u>
β 2	PCx49(1) ^d	~2 Kb <u>EcoRI</u>

20

^aWada et al., supra

^bBoulter et al., (1986), supra

^cGoldman et al., (1987), Cell 48:965-973

^dBoulter et al., (1987), supra

25 * This insert was placed into unmodified pSV2dhfr.

The pSV2dhfr plasmid was modified by first destroying the unique EcoRI site, then inserting a HindIII-EcoRV-EcoRI polylinker between the SV40 early promoter and the dhfr gene. To accomplish this, pSV2dhfr was cut with EcoRI, Klenow-treated, and religated. The resultant plasmid was called pSV2dhfr Δ RI. Plasmid pSV2dhfr Δ RI was digested with HindIII and 0.1 μ g was ligated with a 100:1 molar ratio of unkinased double-stranded oligonucleotide of the following sequence:

23

-23-

AGC TTT CGA TAT CAG AAT TCG¹ (SEQ ID NO: 11)
 AA ~~EST~~ ATA GTC TTA AGCTCGA (SEQ ID NO: 12)
¹
 HindIII EcoRV EcoRI destroyed HindIII

5 The ligation reaction was transformed into MC1061 bacterial cells, amp^R colonies were selected, and plasmid was isolated. Correctly modified plasmid demonstrated a 350 bp band upon digestion with PvuII/EcoRI, and was called pSV2+Ldhfr.

10 To create the $\alpha 2$, $\alpha 3$, and $\beta 2$ expression plasmids, 0.1 μ g of pSV2+Ldhfr, or pSV2dhfr in the case of $\alpha 4$, and 0.1 μ g of the subunit specific gel-isolated insert fragment were ligated, and the individual ligations were separately transformed into MC1061 cells.

15 (The parent plasmids were digested with the appropriate enzyme to allow insertion of the insert noted above into the polylinker site prior to ligation.) Amp^R colonies were selected and plasmid was isolated. The final plasmid names and diagnostic bands indicative of the correct
 20 orientation were:

<u>Subunit</u>	<u>Plasmid name</u>	<u>Diagnostic fragments</u>
$\alpha 2$	pSV2dhfr $\alpha 2$	1600 bp <u>BglII</u>
$\alpha 3$	pSV2dhfr $\alpha 3$	600 bp <u>PvuII</u> ; 850 bp <u>BamHI</u>
$\alpha 4$	pSV2dhfr $\alpha 4$	800 bp <u>PvuII/SstI</u>
$\beta 2$	pSV2dhfr $\beta 2$	1800 bp <u>PvuII</u>

These final plasmids have the subunit insert placed in functional association downstream of the SV40 early
 30 promoter.

C. TK⁺ Selection Plasmids

The TK⁺ plasmid cotransfected into Ltk⁺
 35 cells along with the nNACHR subunit-expressing plasmids

24

-24-

was either pThx59 [Zipser, et al., Proc. Natl. Acad. Sci. 78:6276-6280 (1981)] which encodes the wildtype TK gene, or pThx24 (ibid.) which encodes a crippled TK gene.

5 D. Reporter Gene Expression Plasmid

A reporter gene expression plasmid comprised of the CAT gene regulated by the c-fos promoter, plasmid pFC4 [(Deschamps et al., Science 10 230:1174-1177 (1985)], can also be cotransfected into the cells.

E. Transfection and TK⁺ Selection

15 The CaPO₄ transfection procedure used in the development of the rat nNACHR-expressing cell lines was that of Wigler, et al. (1979), Proc. Natl. Acad. Sci. 76:1373-1376.

Briefly, Ltk⁺ cells were grown in
20 nonselective medium [D + 10 (Dulbecco's modified Eagle's medium + 10% calf serum), 100 U/ml penicillin, and 100 µg/ml streptomycin] in a 10 cm-sized dish, to 20% confluence. The three circular vector DNAs were precipitated with CaPO₄ and added to the cell monolayer.
25 The vector concentrations were as follows:

Thx24:α _x β ₂	2µg:2µg:2µg/ml
Thx59:α _x β ₂	0.25µg:2µg:2µg/ml

30 The transfected cells were allowed to grow for two days in nonselective medium. After two days, the cells were passed and non-selective media was replaced with selective HAT medium (D + 10 + 15 µg/ml hypoxanthine + 1
35 µg/ml aminopterin + 5 µg/ml thymidine), and the cells

25

-25-

were left to grow for 10-15 days, during which time the cells were "fed" fresh selective (HAT) medium every 3-4 days. After 10-15 days, colonies appeared which indicated acceptance and expression of at least the plasmid carrying the TK gene. Colonies were transferred into separate wells of a 24-well dish and grown in selective medium for seven days, at which time individual colonies were passed into 6-well dishes and grown for another seven days in selective medium. To provide cells for freezing and subsequent molecular and functional receptor analyses, the individual clones in the 6-well dishes were passed to 100 ml dishes in selective medium for 5-7 days.

Example 3

CHARACTERIZATION OF CELL LINES EXPRESSING NACHR

The cell lines developed according to the methods of Example 2 were characterized using one or more of the methods described below.

A. Northern or slot blot analysis for expression of α - and β - subunit encoding messages

Total RNA was isolated from 1×10^7 cells and 10-15 μ g of RNA from each cell type were used for Northern or slot blot hybridization analysis. The inserts from the rat nNACHR-encoding plasmids were nick-translated and used as probe. In addition, the β -actin gene sequence [(Cleveland et al., Cell 20:95-105 (1980))] was nick-translated and used as a control probe on duplicate filters to confirm the presence or absence of RNA on each blot and to provide a rough standard for use in quantitating differences in α - or β -specific mRNA

26

-26-

levels between cell lines. The Northern and slot blot hybridization and wash conditions were as follows:

Hybridization: 5XSSPE, 5X Denhardts,
50% formamide, 42°C
Wash: 0.2XSSPE, 0.1% SDS, 65°C

The results of these analyses showed that, while the amount of counts per minute corresponding to actin message was fairly constant among the various cells lines, the levels of α - and β -specific messages varied. Cell lines testing positive for both α - and β -specific mRNA were further tested for functional receptors.

B. Nicotine-binding assay

Cell lines which demonstrated α - and β -specific mRNA were analyzed for their ability to bind nicotine, as compared to three control cell lines: the neuronally-derived cell lines PC12 (Boulter *et al.*, (1986), *supra*) and IMR32 (Clementi, *et al.* (1986); *Int. J. Neurochem* 47:291-297, and the muscle-derived cell line BC3H1 (Patrick, *et al.*, (1977); *J. Biol. Chem.* 252:2143-2153). The assay was conducted as follows:

Just prior to being assayed, the transfected cells were removed from plates by scraping. PC12, BC3H1, and IMR32 (which had been starved for fresh media for seven days). Control cell lines were removed by rinsing in 37°C assay buffer (50mM Tris/HCl, 1 mM MgCl₂, 2 mM CaCl₂, 120 mM NaCl, 3 mM EDTA, 2 mg/ml BSA and 0.1 % aprotinin at pH7.4). The cells were washed and resuspended to a concentration of $1 \times 10^6/250 \mu\text{l}$. To each plastic assay tube was added 250 μl of the cell solution, 15 nM ³H-nicotine, with or without 1 mM cold nicotine, and assay buffer to make a final volume of 500 μl . The

27

-27-

assays for the transfected cell lines were incubated for 30 min at room temperature; the assays of the positive control cells were incubated for 2 min at 1°C. After the appropriate incubation time, 450 µl aliquots of assay volume were filtered through Whatman GF/C glass fiber filters which had been pretreated by incubation in 0.05% polyethyleneimine for 24 hours at 4°C. The filters were then washed twice, with 4 ml each wash, with ice cold assay buffer. After washing, the filters were dried, added to vials containing 5 ml scintillation fluid and then counted.

The IC₅₀ values for displacement of specifically bound ³H-nicotine in the three control cell lines were:

T280X

Cell line	Nicotine concentration required to displace 50% bound nicotine (IC ₅₀)
BC3H1	90 µM
PC12	40 µM
IMR32	35 µM

C. ⁸⁶Rb ion-flux assay

The ability of nicotine or nicotine agonists and antagonists to mediate the influx of ⁸⁶Rb into transfected and control cells has been found to provide an indication of the presence of functional NACHRs on the cell surface. The ⁸⁶Rb ion-flux assay was conducted as follows:

1. The night before the experiment, the cells were plated at 2 x 10⁶ per well (i.e., 2 ml per well) in a 6-well polylysine-coated plate.

28

-28-

2. The culture medium was decanted and the plate was washed with 2 ml of assay buffer (50 mM hepes, 260 mM sucrose, 5.4 mM KCl, 1.8 mM CaCl₂, 0.8 mM MgSO₄, 5.5 mM glucose) at room temperature.

5

3. The assay buffer was decanted and 1 ml of assay buffer, containing 2 μ Ci/ml ⁸⁶Rb, with 5 mM ouabain and agonist or antagonist in a concentration to effect a maximum response, was added.

10

4. The plate was incubated on ice at 1°C for 4 min.

5. The buffer was decanted into a waste container and each well was washed with 3 ml of assay buffer, followed by two washes of 2 ml each.

6. The cells were lysed with 2 x 0.5 ml of 0.2% SDS per well and transferred to a scintillation vial containing 5 ml of scintillation fluid.

20

7. The vials are counted and the data calculated.

25 The positive control cells provided the following data in this assay:

29

-29-

	<u>Agonist</u>	<u>PC12</u>		<u>IMR32</u>	
		<u>EC₅₀</u>	<u>Maximum response</u>	<u>EC₅₀</u>	<u>Maximum response</u>
5	nicotine	52 μ M	2.1X ^a	18 μ M	7.7X ^a
	carbamylcholine (CCh)	35 μ M	3.3X ^b	230 μ M	7.6X ^c
	cytisine	57 μ M	3.6X ^d	14 μ M	10X ^e
	<u>Antagonist</u>				
10	d-tubocurarine	0.81 μ M		2.5 μ M	
	mecamylamine	0.42 μ M		0.11 μ M	
	hexamethonium	nd ^f		22 μ M	
	atropine	12.5 μ M		43 μ M	

15

^a 200 μ M nicotine
^b 300 μ M CCh
^c 3mM CCh
^d 1mM cytisine
^e 100 μ M cytisine
^f nd=not determined

20

D. Nicotine-induced c-fos promoted expression of CAT

25

In cell lines developed by cotransfection of the pFC4 c-fos-CAT plasmid along with the nNACHR subunit-encoding plasmids and the marker plasmid, the functionality of the nNACHRs can be indirectly evaluated by measuring the level of CAT activity. The CAT activity assay can be performed by any of the known methods in the art. See, for example, Nielsen et al., Anal. Biochem. 179:19-23 (1989).

35

E. Xenopus oocytes assay

30

-30-

The functionality of the nNACHR expressed in transfected cells or encoded by the human neuronal NACHR subunit-encoding cDNAs can be evaluated in the Xenopus oocytes system. See Dascal, N. (1987), CRC Crit. Rev. Biochem. 22:317-387, for a review of the use of Xenopus oocytes to study ion channels. RNA from transfectant cell lines or transcribed in vitro from the subunit-encoding cDNAs is injected into oocytes for translation into functional protein. The function of the expressed nNACHR can be assessed by a variety of electrophysiological techniques, including intracellular voltage recording, two-electrode voltage clamp, and patch clamp methods. The cation-conducting channel intrinsic to the NACHR opens in response to acetylcholine (ACh) or other nicotinic cholinergic agonists, permitting the flow of transmembrane current carried predominantly by sodium and potassium ions under physiological conditions. This current can be monitored directly by the voltage clamp techniques, or indirectly by intracellular voltage recording, wherein changes in membrane potential due to the net entry of cations are measured. With the intracellular voltage method, perhaps the simplest technique, a depolarization is recorded upon external application of agonist, signifying the presence of functional receptors in the oocyte membrane.

In a typical experiment to evaluate the functionality of nNACHR subunit-encoding transcripts, 15 oocytes were injected with 5 ng of a 1:1 mixture of an α and a β transcript. Other oocytes were injected with water to serve as negative controls. The oocytes were then incubated at 19°C for 2-5 days in OR-2, an oocyte Ringer's solution of the following composition (concentration in mM): NaCl, 82.5; KCl, 2.5; Na₂HPO₄, 1; HEPES, 5; CaCl₂, 1; MgCl₂, 1; pH = 7.8. For electrophysiological recording, OR-2 of identical

31

-31-

composition except at pH = 7.5 was used as the basis of drug-containing solutions of the bath and agonist application pipet. During continuous intracellular voltage recording in a bath of OR-2 containing 1 μ M atropine to block endogenous muscarinic acetylcholine receptor responses, a pipet containing 100 μ M ACh was used to intermittently apply ACh by a local perfusion method in which the ACh is diluted by a factor of about 3-10 upon application to the oocyte.

Healthy oocytes have resting potentials in the range of -50 to -70 mV. Depolarizations due to ACh ranged from several mV to about 30 mV in different batches of oocytes injected with NACHR subunit-encoding transcripts. (Responses within a given batch of oocytes tended to be of similar magnitude.) The depolarizing responses to ACh were reversibly blocked by 100 μ M d-tubocurarine, added to the bath. By contrast, water-injected oocytes did not respond at all to ACh administration under these conditions.

In a typical experiment to evaluate the nNACHR subunit-encoding RNA from transfected cell lines, total RNA was isolated from the cells and 50 ng were injected into oocytes. The oocytes were incubated and treated with acetylcholine, atropine, and d-tubocurarine as described above. Negative control oocytes were injected with RNA from a negative control cell line transfected with parent plasmid lacking a nNACHR subunit-encoding insert.

Oocytes injected with message from nNACHR-transfected cells demonstrated depolarization when treated with acetylcholine. The depolarization was blocked with d-tubocurarine. The negative control oocytes were unresponsive, as expected.

Alternatively, the functionality of nNACHRs expressed in transfected cells can be studied by standard

32

-32-

electrophysiological techniques such as intracellular voltage recording or patch clamp, analogous to the methods described for oocytes.

5

Example 4

CELL LINES EXPRESSING FUNCTIONAL nNACHRs

Several cell lines were generated employing the procedures of Example 2. The resulting cell lines were then analyzed employing the assay methods described in Example 3. Results for several newly prepared cell clones are summarized below:

T330X

	<u>Cell line</u>	<u>Subunits</u>	<u>RNA analysis</u>	<u>Binding</u>	<u>Rb flux</u>	<u>Oocytes</u>
15	592F	$\alpha 2\beta 2^a$	+/+ ^b	+ ^c	nd	+ ^d
	243C	$\alpha 3\beta 2$	+/+	+	nd	+
	244A	$\alpha 4\beta 2$	+/+	+	nd	+
	244I	$\alpha 4\beta 2$	+/+	+	nd	nd

20

nd=not determined

^a subunits are from rat nNACHR^b +/+ indicates that α - and β -specific mRNA was detected^c + indicates that the cell line binds agonist in a manner similar to positive control cells25 ^d + indicates that ACh induces membrane depolarization which was blocked by d-tubocurarine.

These results show that functional nNACHRs are expressed by mammalian cells transfected with DNA encoding an α -subunit and a β -subunit of the nNACHR.

The invention has been described in detail with reference to certain particular embodiments thereof. It will be understood, however, that variations and modifications can be effected within the spirit and scope of the invention.

35

add B'

33